

**Osaka University of Economics Working Paper
Series
No. 2009-6**

Strong Convexity of NTU Games

**Takuya Masuzawa
Faculty of Economics, Osaka University of Economics
December 8, 2009**

Strong Convexity of NTU Games

Takuya Masuzawa *

December 8, 2009

Abstract

In this paper, we provide an alternative definition of NTU convexity, *strongly ordinal convexity*. We show that if a game is strongly ordinal convex, then any marginal worth vector is in the core, and any marginal contribution is increasing. Some economic examples satisfy this convexity.

Keywords Cooperative game; Convex game; NTU game; Core; Super-modularity

JEL classification codes: C62; D52; D53

1 Introduction

The convexity of NTU game was first defined as an extension of the TU convex game by Vilkov (1977), called *weakly ordinal convexity* in this paper. While weakly ordinal convex games arise in various economic applications¹, they do not inherit interesting properties from the TU convex game, any of which also characterizes the TU convexity (see Sharkey, 1981; Hendrickx et al. 2000, 2002). Moreover, these properties do not imply the weakly ordinal convexity. In this paper, I propose a strong concept of NTU convexity, *strongly ordinal convexity*, to inherit the various properties of the TU convex games. Roughly speaking, a game is strongly ordinal convexity iff for all coalition T , the proper contribution of coalition $S \subset N \setminus T$ in $S \cup T$ is super additive with respect to S . We show that a strongly ordinal convex game is weakly ordinal convex, any marginal contribution of this game is increasing, and its core comprises all marginal worth

*Faculty of Economics, Osaka University of Economics, 2-2-8 Osumi, Higashi-yodogawa-ku, Osaka-shi. Email: masuzawa@osaka-ue.ac.jp

¹See Peleg (1984).

vectors. Further, we show that some economic examples are strongly ordinal convex.

2 Results

2.1 Preliminary

Let $\mathbb{R}$ be the set of real numbers. Let $N = \{1, 2, \dots, n\}$ be a finite set of players. A nonempty subset of N is called a coalition. A payoff vector is an element in $\mathbb{R}^N$. For all $S \in 2^N$ and $x \in \mathbb{R}^N$, x^S denotes the projection of x to $\mathbb{R}^S$, and $x(S)$ denotes $\sum_{i \in S} x_i$. For all $x, y \in \mathbb{R}^N$, we write $x^S \geq y^S$ (resp. $x^S \gg y^S$) iff $x^i \geq y^i$ (resp. $x^i > y^i$) for all $i \in S$. A *coalitional form game* specifies a set of payoffs that can be obtained by coalition S by itself. A *TU coalitional form game*, v , is a function from 2^N to $\mathbb{R}$ with $v(\emptyset) = 0$. A payoff $x \in \mathbb{R}^N$ can be obtained by S by itself iff $x(S) \leq v(S)$. A *core* of v is a set of payoff vectors $x \in \mathbb{R}^N$ such that $x(S) \geq v(S)$ for all $S \in 2^N$ and $x(N) \leq v(N)$. A game v is *convex* if $v(S) + v(T) \leq v(S \cap T) + v(S \cup T)$ for all $S, T \in 2^N$. This definition is equivalent to the property, called *the increasing marginal contribution*, that $v(Q \cup R) - v(Q) \leq v(T \cup R) - v(T)$ for all $Q \subseteq T \subseteq N \setminus R$. Let σ be a permutation of N . The *marginal worth vector* for σ is a payoff vector, denoted by m_σ , such that $m_\sigma^{\sigma(1)} = v(\sigma(1))$; for all $k > 1$, $m_\sigma^{\sigma(k)} = v(\sigma(\{1, 2, \dots, k\})) - v(\sigma(\{1, 2, \dots, k-1\}))$. Shapley (1971) and Ichiishi (1981) showed that a TU game is convex iff any marginal worth vector is in the core.

An *NTU game* is a correspondence from 2^N to $\mathbb{R}^N$ such that for all $x, y \in \mathbb{R}^N$, if $x^S \geq y^S$ and $x \in V(S)$, then $y \in V(S)$, and that $V(\emptyset) = \emptyset$. Any situation described by a TU game v is also expressible by an NTU game such that $V_v(S) := \{x \in \mathbb{R}^N \mid v(S) \geq x(S)\}$. For all $S \in 2^N$, let $D(S)$ denote the set of payoff vectors x such that there exists no $y \in V(S)$ such that $y^S \gg x^S$. Let $\hat{V}(S)$, and $\hat{D}(S)$ denote the projections of $V(S)$ and $D(S)$ to $\mathbb{R}^S$ respectively. A core of V is defined by $V(N) \cap (\cap_{S \subseteq N} D(S))$.

An NTU game is *weakly ordinal convex* if for all $S, T \in 2^N \setminus \{\emptyset\}$, $V(S) \cap$

$V(T) \subseteq V(S \cup T) \cup V(S \cap T)$. This notion is an extension of the TU convexity to NTU game in the sense that V_v is weakly ordinal convex iff v is convex. The weakly ordinal convexity plays important roles in the effectivity function analyses (see chapter 6 of Peleg 1984). Further, the core of weakly ordinal convex game is nonempty and coincides with the unique vNM stable set (see Greenberg, 1985; Peleg, 1986).

2.2 Strong Convexity

For all $R \in 2^N \setminus \{\emptyset\}$ and all $Q \subseteq N \setminus R$, x^R is a *proper contribution* of R to x^Q iff $x \in V(R \cup Q) \cap D(Q)$. That is, x^R is a payoff that can be allocated to R if R collaborates with Q and gives x^Q to Q , which cannot be improved upon by Q by itself. Now, we define our main concept:

Definition 1 *An NTU game is strongly ordinal convex if*

$$V(S) \cap D(S \cap T) \cap V(T) \subseteq V(T \cup S) \text{ for all } S, T \in 2^N.$$

By taking $R_0 := S \cap T$, $R_1 := S \setminus T$, and $R_2 := T \setminus S$, one can easily see that a game is strongly ordinal convex iff proper contributions are super-additive with respect to coalitions: for all $R_0, R_1, R_2 \subseteq N$ such that $R_i \cap R_j = \emptyset$ for all $i \neq j$, if both of x^{R_1} and x^{R_2} are proper contributions of R_1 and R_2 to x^{R_0} respectively, then (x^{R_1}, x^{R_2}) is a proper contribution of $(R_1 \cup R_2)$ to x^{R_0} . This condition was first proposed by Milgrom and Shannon (1996), who attempted to define a general notion of NTU convexity².

First, we show that this definition is an alternative extension of the TU convexity.

Theorem 2 *For any TU game v , an NTU game V_v defined by $V_v(S) := \{x \in \mathbb{R}^N \mid v(S) \geq x(S)\}$ for all $S \in 2^N \setminus \{\emptyset\}$ is strongly ordinal convex iff v is convex.*

²Their generalization is not successful as Theorem 8 of their paper is mathematically incorrect.

Proof. For all $x \in \mathbb{R}^N$, $x \in V_v(S) \cap D_v(S \cap T) \cap V_v(T)$ iff $-x(S \cap T) \leq -v(S \cap T)$, $x(S) \leq v(S)$, and $x(T) \leq v(T)$. Then, for all $x \in V_v(S) \cap D_v(S \cap T) \cap V_v(T)$,

$$v(S) + v(T) - v(S \cap T) \geq x(S \cup T). \quad (1)$$

If v is convex, then the left hand side of (1) is smaller than or equal to $v(S \cup T)$, which implies that $x \in V_v(S \cup T)$. Conversely, suppose that V_v is strongly ordinal convex. We can choose $x \in V_v(S) \cap D_v(S \cap T) \cap V_v(T)$ to satisfy the equality of (1). The right hand side of (1) is then smaller than or equal to $v(S \cup T)$. It follows that v is convex. $\square$

Our convexity implies the convexity mentioned in Vilkov (1977).

Theorem 3 *A strongly ordinal NTU game is weakly ordinal convex.*

Proof. Suppose that V is strongly ordinal convex. Let $x \in V(S) \cap V(T)$. If $x \in D(S \cap T)$, then we have $x \in V(S \cup T)$. On the other hand, if $x \notin D(S \cap T)$, then there exists $y \in V(S \cap T)$ such that $y^{S \cap T} \gg x^{S \cap T}$ and, therefore, $x^{S \cap T}$ is also in $\hat{V}(S \cap T)$. $\square$

The *marginal worth vector* for σ , m_σ , is defined by $m_\sigma^{\sigma(k)} := \max\{y^{\sigma(k)} \mid (m_\sigma^{\sigma(\{1,2,\dots,k-1\})}, y^{\sigma(k)}) \in \hat{V}(\sigma(\{1,2,\dots,k\}))\}$. Hendrickx et al. (2000, 2002) illustrated that the core of an ordinal convex game does not necessarily comprise all marginal worth vectors. On the other hand, we have the following:

Theorem 4 *The core of a strongly ordinal NTU game comprises all marginal worth vectors.*

Proof. Consider the induction on $|N|$. Let σ be a permutation and W be a restriction of V to 2^T , where $T := N \setminus \{\sigma(n)\}$. Suppose that there exist $S \neq \emptyset$ and $y \in \mathbb{R}^N$ such that $y \in V(S)$ and $y^S \gg m_\sigma^S$. By the induction hypothesis, m_σ^T is in the core of W . Thus, $\sigma(n) \in S$ and $m_\sigma \in D(S \cap T)$. Then, $(m_\sigma^T, y^{\sigma(n)}) \in V(S) \cap D(S \cap T) \cap V(T)$. From strongly ordinal convexity, $(m_\sigma^T, y^{\sigma(n)}) \in V(N)$; this contradicts the definition of $m_\sigma^{\sigma(n)}$. $\square$

Economic Examples Economic models in Masuzawa (2004) satisfy the condition that $V(S) \cap V(T) \subseteq V(S \cup T)$ for all $S, T \in 2^N$. Obviously, they are strongly ordinal convex. Note that this condition is not adequate for an alternative concept of NTU convexity since V_v does not satisfy this condition for all TU games v .

Weakly ordinal convex games arise in voting theory³. On the other hand, we have the following example.

Example 5 Consider a three-person majority voting game with two alternatives, a and b . Suppose that a and b give the players $(x^1, x^2, x^3) = (1, 0, 0)$ and $(y^1, y^2, y^3) = (0, 1, 0)$ respectively. Then,

$$V(\{i\}) = \{x \mid x^i \leq 0\} \text{ for } i = 1, 2, 3;$$

$$V(\{i, 3\}) = \{x \mid (x^i, x^3) \leq (1, 0)\} \text{ for } i = 1, 2;$$

$$V(\{1, 2\}) = \{x \mid \max\{x^1, x^2\} \leq 1 \text{ and } \min\{x^1, x^2\} \leq 0\};$$

$$V(\{1, 2, 3\}) = \{x \mid x^3 \leq 0 \text{ and } x \in V(\{1, 2\})\}.$$

This is an weakly ordinal convex game that is not strongly ordinal convex.

2.3 Increasing marginal contribution

For all $R \subseteq N$ and $Q \subseteq N \setminus R$, x^R is a *marginal contribution* to x^Q if it is a proper contribution of R to x^Q and $x^Q \in \hat{V}(Q)$. I show that any marginal contribution of a strongly ordinal convex game is increasing in Q . By taking $Q := S \cap T$ and $R := S \setminus Q$, we can see that an NTU game is strongly ordinal convex iff $V(R \cup Q) \cap D(Q) \cap V(T) \subseteq V(T \cup R)$ for all $Q \subseteq T \subseteq N \setminus R$. It follows that for all $Q \subseteq T \subseteq N \setminus R$,

$$V(R \cup Q) \cap (D(Q) \cap V(Q)) \cap (D(T) \cap V(T)) \subseteq V(T \cup R). \quad (2)$$

Condition (2) says that if x^R is a marginal contribution to x^Q , then it is also a marginal contribution to all $y^T \in \hat{D}(T) \cap \hat{V}(T)$ such that $y^Q = x^Q$. Note that for any TU game v , V_v satisfies (2) iff it satisfies the increasing marginal contribution property: $v(Q \cup R) - v(Q) \leq v(T \cup R) - v(T)$.

³See chapter 6 of Peleg (1984).

Note that in some weakly ordinal convexity games, a marginal contributions are not increasing in our sense.

Example 6 Consider a three-person game defined by

$$V(\{i\}) = \{x \mid x^i \leq 0\} \text{ for } i = 1, 2, 3;$$

$$V(\{1, 2\}) = \{x \mid x^1 + x^2 \leq 1\};$$

$$V(\{i, 3\}) = \{x \mid (x^i, x^3) \leq (0, 1)\} \text{ for } i = 1, 2;$$

$$V(\{1, 2, 3\}) = \{x \mid x^1 + x^2 + x^3 \leq 1\}.$$

This game is weakly ordinal convex. However, the marginal contribution of $\{3\}$ to $x^1 = 0$ is 1 while that of $\{3\}$ to $y^{\{1,2\}} \in \hat{V}(\{1, 2\}) \cap \hat{D}(\{1, 2\})$ is 0.

A payoff vector a^R is *acceptable* for $Q \subseteq N \setminus R$ iff it is a proper contribution to some a^Q . Milgrom and Shannon (1996) introduced the following property, called *the scale merit of acceptability* in this paper: for all $Q \subseteq T \subseteq N \setminus R$ and all $a^R \in \mathfrak{R}^R$, if a^R is acceptable for Q , then a^R is also acceptable for T .

Theorem 7 *Let V be an NTU game. Assume that for all $S \in 2^N \setminus \{\emptyset\}$, $V(S)$ is closed and nonempty, and that for all $b \in V(S)$, $\{x^S \in \hat{V}(S) \mid x^S \geq b^S\}$ is bounded in $\mathfrak{R}^S$. If V is strongly ordinal convex, then V has the scale merit of acceptability.*

Proof. It suffices to consider the case where $Q \subsetneq T = N \setminus R$. Suppose that a^R is acceptable for Q . Then, there exists $a^Q \in \hat{D}(Q)$ such that $(a^R, a^Q) \in \hat{V}(Q \cup R)$. Since $V(Q)$ is nonempty, there exists $x^Q \in \hat{V}(Q)$ such that $x^Q \leq a^Q$. Since $V(Q)$ is closed, there exists $b^Q \in \hat{D}(Q) \cap \hat{V}(Q)$ such that $(a^R, b^Q) \in \hat{V}(R \cup Q)$. From strongly ordinal convexity and $V(T \setminus Q) \neq \emptyset$, there exists $y^{T \setminus Q} \in \mathfrak{R}^{T \setminus Q}$ such that $(y^{T \setminus Q}, b^Q) \in \hat{V}(T)$. Since $\{x^T \in \hat{V}(T) \mid x^T \geq y^T\}$ is bounded and closed, we can choose $c^{T \setminus Q}$ such that $(b^Q, c^{T \setminus Q}) \in \hat{V}(T) \cap \hat{D}(T)$. Then, $(a^R, b^Q, c^{T \setminus Q}) \in V(R \cup Q) \cap D(Q) \cap V(T)$. From strongly ordinal convexity, $(a^R, b^Q, c^{T \setminus Q}) \in V(T \cup R)$. Further, since $(a^R, b^Q, c^{T \setminus Q}) \in D(T)$, a^R is also acceptable for T . $\square$

The converse of Theorem 7 is, however, not true.

Example 8 Consider a three-person game such that

$$V(\{i\}) = \{x \mid x^i \leq 0\} \text{ for all } i = 1, 2, 3;$$

$$V(S) = \{x \mid \min_{i \in S} \{x^i\} \leq 1 \text{ and } \max_{i \in S} \{x^i\} \leq 2\} \text{ if } |S| = 2;$$

$$V(S) = \{x \mid x(S) \leq 3 \text{ and } \min_{i \in S} \{x^i\} \leq 0\} \text{ if } |S| = 3.$$

This game satisfies the scale merit of acceptability. The core of the game is, however, empty.

Hendrickx et al.(2000, 2002) proposed *the coalition-merge property*: for all $Q \subseteq T \subseteq N \setminus R$, all $a \in D(Q) \cap V(Q \cup R) \cap (\cap_{i \in Q} D(\{i\}))$, and all $b \in V(T)$, $(a^R, b^{N \setminus R}) \in V(T \cup R)$. This requirement is strong in that a^R is compatible with all $b^T \in \hat{V}(T)$, which is chosen independently of a^Q . On the other hand, note that, in our definition, b^T is assumed to be an extension of a^Q .

Example 9 Consider a four-person game defined by

$$V(\{1, 2, 3, 4\}) = \{x \mid x(\{1, 2, 3, 4\}) \leq 2\};$$

$$V(\{1, 2, 3\}) = \{x \mid (x^1, x^2, x^3) \leq (1, 0, 1)\};$$

$$V(\{1, 2, 4\}) = \{x \mid (x^1, x^2, x^4) \leq (0, 1, 1)\};$$

$$V(S) = \{x \mid x^i \leq 0 \text{ for all } i \in S\} \text{ otherwise.}$$

This game does not satisfy the coalition-merge property. Let $Q = \{1, 2\}$, $R = \{3\}$, $T = \{1, 2, 4\}$. Then, $(a^1, a^2, a^3, x) = (1, 0, 1, x) \in D(Q) \cap V(Q \cup R) \cap (\cap_{i \in Q} D(\{i\}))$ and $(b^1, b^2, b^3, b^4) = (0, 1, 0, 1) \in V(T)$. However, $(a^R, b^T) = (b^1, b^2, a^3, b^4) = (0, 1, 1, 1) \notin V(T \cup R)$. On the other hand, it is strongly ordinal convex.

The following example from Hendrickx et al. (2000, example 4.6) also illustrates that the coalition-merge property does not necessarily mean the properties discussed in this paper:

Example 10 Consider a four-person game defined by

$$V(\{1, 2, 3, 4\}) = \{x \mid x(\{1, 2, 3, 4\}) \leq 7\};$$

$$V(S) = \{x \mid x(S) \leq 4\} \text{ if } |S| = 3;$$

$$V(S) = \{x \mid \max_{i \in S} x^i \leq 1\} \text{ if } |S| = 2;$$

$$V(S) = \{x \mid x^i \leq 0 \text{ for all } i \in S\} \text{ otherwise.}$$

Hendrickx et al. (2000) showed that this game has the coalition-merge property but is not weakly ordinal convex. It does not satisfy the scale merit of acceptability. To see this, consider $x^1 = 4$, which is a proper contribution of player 1 to $(x^2, x^3) = (1, -1)$. It is not, however, a proper contribution of player 1 to any (y^2, y^3, y^4) , which must be smaller than 3. It follows that the coalition merge property does not imply strong convexity. Further, $(x^1, x^2, x^3, x^4) = (4, 1, -1, 4) \in V(\{1, 2, 3\}) \cap V(\{2, 3\}) \cap D(\{2, 3\}) \cap V(\{2, 3, 4\}) \cap D(\{2, 3, 4\})$ while $(x^1, x^2, x^3, x^4) \notin V(\{1, 2, 3, 4\})$. Thus, a marginal contribution is not increasing in our sense.

3 Concluding Remarks

We defined strongly ordinal convexity as an extension of the TU convexity to NTU game, which is a subclass of the weakly ordinal convex games of Vilkov (1977). Further, any marginal contribution of this game is increasing and the core of this game comprises all marginal worth vectors.

Reference

- Greenberg, Joseph(1985), Cores of convex games without side payments. *Mathematics of Operations Research* 10: 523-525.
- Hendrickx, Ruud, and Timmer, Judith, and Borm, Peter (2000), On convexity for NTU-games, Tilburg University, CentER Working Paper No. 2000-108. <http://ssrn.com/abstract=250792> or doi:10.2139/ssrn.250792.
- Hendrickx, Ruud, and Timmer, Judith, and Borm, Peter (2002), A note on NTU convexity, *International Journal of Game Theory* 31: 29-37
- Ichiishi, Tatsuro (1981), Super-modularity: applications to convex games and to the greedy algorithm for LP, *Journal of Economic Theory* 25: 283-286.
- Masuzawa, Takuya (2004) Punishment strategies make the α -coalitional game ordinally convex and balanced, *International Journal of Game Theory* 32:

479-483.

- Milgrom, Paul and Cris Shannon (1996), Generalized convex games,
<http://elsa.berkeley.edu/users/cshannon/wp/core.pdf>, Berkley University
- Peleg, Bezalel (1984), *Game Theoretic Analysis of Voting in Committees*, Cambridge University Press, Cambridge.
- Peleg, Bezalel (1986), A Proof that the core of an ordinal convex game is a von Neumann-Morgenstern colution, *Mathematical Social Sciences* 11: 83-87.
- Shapley, Lloyd (1971), Cores of convex games, *International Journal of Game Theory* 1: 11-26.
- Sharkey, William (1981), Convex games without side payments, *International Journal of Game Theory* 10: 101-6.
- Vilkov, V.B. (1977), Convex games without side payments, *Vestnik Leningradskiva University*, 7: 21-24 (in Russian).